

Anticommutativity Equation in Topological Quantum Mechanics

Vyacheslav Lysov[†]

Institute of Theoretical and Experimental Physics, 117259, Moscow, Russia
and
Moscow Institute of Physics and Technology, 141700, Moscow, Russia

Abstract

We consider topological quantum mechanics as an example of topological field theory and show that its special properties lead to numerous interesting relations for topological corellators in this theory. We prove that the generating function $\mathcal{F}$ for thus corellators satisfies the anticommutativity equation $(\mathcal{D} - \mathcal{F})^2 = 0$. We show that the commutativity equation $[dB, dB] = 0$ could be considered as a special case of the anticommutativity equation.

1. During the last two decades there has been much interest to quantum field theories whose special corellators do not depend on coordinates and metric. These theories are called topological[1]. The most celebrated examples include Chern-Simons theory [2], $N = 2$ twisted gauge theories[3], topological sigma models[4]. Here we consider yet another, much simpler example of topological theory which is a subsector of supersymmetric quantum mechanics. We call it topological quantum mechanics.

An explicit construction of many topological theories is given by a BRST-like symmetry operator Q , $Q^2 = 0$ so that the energy momentum tensor of the theory has the special form [3]

$$T_{\mu\nu} = \{Q, G_{\mu\nu}\} \quad (1)$$

where $G_{\mu\nu}$ is some tensor. This formula leads to many interesting corollaries. In particular, it makes corellators of Q -closed operators independent of metric and coordinates, that is topologically invariant.

In the case of one dimensional theory, the energy momentum tensor has only one component and is equal to the Hamiltonian

$$H = T_{00} = \{Q, G_{00}\} = \{Q, G_+\} \quad (2)$$

In topological quantum mechanics, the objects of study are corellators $\langle \Phi_{A_1}(t_1) \dots \Phi_{A_n}(t_n) \rangle$ of Q -closed operators. They would naively depend of n times $t_1 < \dots < t_n$. However, since the energy momentum tensor is anticommutator of Q and G_+ , the corellator actually does not depend on t 's and is given by a factorization formula

$$\begin{aligned} \langle \Phi_{A_1}(t_1) \dots \Phi_{A_n}(t_n) \rangle &= \langle 0 | e^{-t_1 H} \Phi_{A_1} e^{(t_1 - t_2) H} \Phi_{A_2} \dots e^{(t_{n-1} - t_n) H} \Phi_{A_n} e^{t_n H} | 0 \rangle = \\ &= \lim_{(t_k - t_{k-1}) \rightarrow \infty} \langle 0 | \Phi_{A_1} e^{(t_1 - t_2) H} \dots e^{(t_{n-1} - t_n) H} \Phi_{A_n} | 0 \rangle = \langle \Phi_{A_1} \rangle \dots \langle \Phi_{A_n} \rangle \end{aligned} \quad (3)$$

[†] e-mail: sllys@gate.itep.ru

It means that it is sufficient to study the corellators of one operator Φ_A in the theory with the deformed supercharge Q

$$Q \rightarrow Q + \sum \Phi_A T_A = Q + \Phi \quad (4)$$

Later we shall see that this one-point corellator is a total derivative of $\mathcal{F}$ on the space of coupling constants T_A

$$\partial_A \mathcal{F} = \langle \Phi_A \rangle_{\text{deformed}} = \langle T \left\{ \Phi_A e^{\int \{\Phi, G_+\} dt} \right\} \rangle \quad (5)$$

where $T\{\dots\}$ stands for the chronological ordering. In what follows, we carefully formulate the theory and realize that the function $\mathcal{F}$ satisfies an interesting quadratic differential equation which we call anticommutativity equation.

2. Thus, we consider the simplest example of topological field theory, topological quantum mechanics. As usual, there is a nilpotent symmetry operator Q and one can introduce the Hamiltonian in the form (2). All operators act on the space V with the following properties

$$\begin{aligned} V &= V_1 \oplus V_0 \\ HV_0 &= 0 \quad , \quad QV_0 = 0 \\ H &> 0 \quad \text{on } V_1 \end{aligned} \quad (6)$$

This physically means that V is a space of states of our system, V_0 is a space of the vacuum states. We assume that the kernel V_0 of the Hamiltonian consists of Q -closed states, and all non-zero energies have strictly positive real part. In this theory we study the corellators of the form (5), which are the vacuum matrix elements, i.e. the operators from $Hom(V_0, V_0)$ and can be written in the following form[‡]

$$\begin{aligned} F_{\alpha\beta}^{(k)} &= \int_0^\infty \dots \int_0^\infty \langle 0_\alpha | \Phi G_+ e^{-t_1 H} \Phi G_+ e^{-t_2 H} \dots \Phi | 0_\beta \rangle dt_1 \dots dt_{k-1} = \\ &= \frac{1}{k!} \langle T \left\{ \Phi(0) \left(\int_{-\infty}^{+\infty} \{G_+, \Phi(t) dt\} \right)^{k-1} \right\} \rangle_{\alpha\beta} \end{aligned} \quad (7)$$

We want to represent this operator from $Hom(V_0, V_0)$ as an operator from $Hom(V, V)$. The way to do this is to insert Π_0 's (projector on to V_0) at the beginning and at the end. After doing this, one finally obtains the object needed

$$\mathcal{F}^{(k)} = \int_0^\infty \dots \int_0^\infty \Pi_0 \Phi G_+ e^{-t_1 H} \Phi G_+ e^{-t_2 H} \dots \Phi \Pi_0 dt_1 \dots dt_{k-1} \quad (8)$$

The $\Phi(0) \equiv \Phi$ is

$$\Phi(0) = \Phi = \sum \Phi_A T_A \quad (9)$$

Our theory is Euclidean, which means that the evolution of Φ in time can be described by the following formula

$$\Phi(t) = e^{-tH} \Phi(0) e^{tH} \quad (10)$$

Π_0 being a projector onto the vacuum states. One can rewrite Π_0 in terms of H

$$\Pi_0 = \lim_{t \rightarrow \infty} e^{-tH} \quad (11)$$

[‡]Here we used the properties $G_+^2 = 0$ and $G_+ V_0 = 0$ that allow one to interpret F as a corellator in topological theory, but they are not necessary for our main result.

It commutes with Q . Therefore,

$$\begin{aligned} Q\Pi_0 &= \Pi_0 Q = 0 \\ < 0_\alpha | Q &= < 0_\alpha | \Pi_0 Q = 0 \end{aligned} \quad (12)$$

Our operators can be both even and odd, therefore, we introduce a superalgebra to describe their properties, our coupling constants T_A being graded too. Their algebra is

$$\{T_A, T_B\}_s = 0 \quad (13)$$

Where $\{.,.\}_s$ stands for supercommutator. In order to get interesting properties of the corellators, let us consider a special set of operators that satisfy the following algebra

$$\begin{aligned} \{\Phi_A, \Phi_B\}_s &= C_{AB}^K \Phi_K \\ \{Q, \Phi_A\}_s &= 0 \end{aligned} \quad (14)$$

3. Algebra of operators Φ generates numerous commutation relations for $\mathcal{F}^{(k)}$

$$\begin{aligned} \{\mathcal{F}^{(1)}, \mathcal{F}^{(1)}\} &= T_A T_B C_{AB}^K \partial_K \mathcal{F}^{(1)} \\ 2\{\mathcal{F}^{(2)}, \mathcal{F}^{(1)}\} &= T_A T_B C_{AB}^K \partial_K \mathcal{F}^{(2)} \\ 2\{\mathcal{F}^{(1)}, \mathcal{F}^{(3)}\} + \{\mathcal{F}^{(2)}, \mathcal{F}^{(2)}\} &= T_A T_B C_{AB}^K \partial_K \mathcal{F}^{(3)} \\ &\dots \end{aligned} \quad (15)$$

which can be written in a compact form in terms of the $Hom(V_0, V_0)$ valued generating function $\mathcal{F}$

$$\mathcal{F} = \sum \mathcal{F}^{(k)} \quad (16)$$

as

$$\{\mathcal{F}, \mathcal{F}\} = 2\mathcal{D}\mathcal{F} \quad (17)$$

Here we introduced the derivative $\mathcal{D} = \frac{1}{2} T_A T_B C_{AB}^K \partial_K$, One can easily obtain all our corellators as derivatives of $\mathcal{F}$. We now prove this equation in the generic case.

Let us introduce an operator-valued differential form of indefinite degree on R

$$U = e^{-tH} + G_+ dt e^{-tH} \quad (18)$$

We can check that

$$dU + \{Q, U\} = 0 \quad (19)$$

One can construct the differential form on R^{k-1} which takes values in $Hom(V_0, V_0)$ and includes $k+1$ Φ 's

$$\omega^{(k-1)} = \Pi_0 \Phi U_1 \dots \Phi U_k \Phi \Pi_0 \quad (20)$$

Our form is d -closed because $Q\Pi_0 = \Pi_0 Q = 0$ and Φ anticommutes with Q

$$d\omega^{(k-1)} = -\Pi_0 \Phi \{U_1, Q\} \dots \Phi U_k \Phi \Pi_0 - \dots - \Pi_0 \Phi U_1 \dots \Phi \{U_k, Q\} \Phi \Pi_0 = 0 \quad (21)$$

One can consider the integral of the degree $k-1$ component of our form over the boundary of some surface and rewrite it as an integral of $d\omega^{(k-1)}$ over this surface by the Stocks theorem. The surface of integration is the boundary of the k -dimensional cube. The result of integration is as follows

$$\int_0^\infty \dots \int_0^\infty \Pi_0 \Phi (\Pi_0 - 1) \Phi G_+ e^{-t_1 H} \dots G_+ e^{-t_k H} \Phi \Pi_0 dt_1 \dots dt_{k-1} + (perm. \text{ of } \Pi_0 - 1) = 0 \quad (22)$$

The terms that include Π_0 come from the commutator $\{\mathcal{F}, \mathcal{F}\}$, while the other terms come from $T_A T_B C_{AB}^K \partial_K \mathcal{F}$. For our superalgebra of operators Φ_A , one can write (super)Jacobi identities in the following form

$$C_{AB}^K C_{KD}^E = 0 \quad , \quad \mathcal{D}^2 = 0 \quad (23)$$

Our anticommutativity equation can be rewritten as the zero curvature equation

$$\{\mathcal{D} - \mathcal{F}, \mathcal{D} - \mathcal{F}\} = 0 \quad (24)$$

4. Now we can illustrate how this equation (24) works in the simplest case. Define in (15) $\mathcal{F}^{(1)} = \sum F_A T_A$. Then

$$\{F_A, F_B\} = C_{AB}^K F_K \quad (25)$$

For odd operators Φ_A 's, this equation is quite nontrivial. One can see this by considering the matrix example. In the simplest case, our space of states V is the four-dimensional vector space. The operators Q and G_+ are 4×4 matrices

$$Q = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad G_+ = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad H = \{Q, G_+\} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (26)$$

V_0 is the subspace of V with first two zero coefficients. The operators Φ_A 's which satisfy commutation relation (15) are as follows

$$\Phi_A = \begin{pmatrix} a & 0 & 0 & 0 \\ * & -a & * & * \\ * & 0 & & F_A \\ * & 0 & & \end{pmatrix} \quad (27)$$

where stars are some numbers. When considering commutation relations on Φ_A , we are interested only in the right bottom block. It is easy to show that in this block there is anticommutator of the two right bottom blocks of Φ_A and Φ_B . It is what we get commuting matrix elements F_A

$$\{\Phi_A, \Phi_B\} = \begin{pmatrix} * & * & * & * \\ * & * & * & * \\ * & * & & \{F_A, F_B\} \\ * & * & & \end{pmatrix} \quad (28)$$

Thus, we observe that this is only these special commutation relations of matrices Φ_A that provide us with the similar algebra for F_A (i.e. for the right bottom blocks).

Now we can explicitly show this for general V . Consider

$$\{F_A, F_B\} = \Pi_0 \Phi_A \Pi_0 \Phi_B \Pi_0 + \Pi_0 \Phi_B \Pi_0 \Phi_A \Pi_0 \quad (29)$$

Using formula (15) for the projector, one arrives at the following formula

$$\Pi_0 = \lim_{t \rightarrow \infty} e^{-tH} = \lim_{t \rightarrow \infty} \sum (-t)^k \frac{H^k}{k!} = 1 + \left(\sum Q G_+ (-t)^k \frac{H^{k-1}}{k!} + \sum (-t)^k G_+ \frac{H^{k-1}}{k!} Q \right) \Big|_{t \rightarrow \infty} \quad (30)$$

An important property of this representation is

$$\begin{aligned}\Pi_0 \Phi_A Q G_+ (-t)^k \frac{H^{k-1}}{k!} \Phi_B \Pi_0 &= 0 \\ \Pi_0 \Phi_A G_+ (-t)^k \frac{H^{k-1}}{k!} Q \Phi_B \Pi_0 &= 0\end{aligned}\tag{31}$$

Hence, the only term surviving in the expression for the projector in (29) is the c -number term in (30) and

$$\{F_A, F_B\} = \Pi_0 \Phi_A \Phi_B \Pi_0 + \Pi_0 \Phi_B \Phi_A \Pi_0 = C_{AB}^K F_K\tag{32}$$

5. Our anticommutativity equation contains a commutativity equation introduced in [5,6,7,8][§] for special operators. The commutativity equation has the form

$$[B_\mu, B_\nu] = 0 \quad , \quad B_\mu = \frac{\partial B}{\partial \tau_\mu}\tag{33}$$

Indeed, consider a new odd operator G_- that satisfies

$$G_-^2 = 0 \quad , \quad \{G_-, G_+\} = 0 \quad , \quad \{G_-, Q\} = 0 \quad , \quad G_- V_0 = 0\tag{34}$$

Consider even operators Φ_μ which satisfy relations (14) with $C = 0$ and

$$[[\Phi_\mu, G_-], \Phi_\nu] = 0\tag{35}$$

Thus properties allow one to add the new odd operators $\Phi'_\mu = [\Phi_\mu, G_-]$ to our algebra because they satisfy (14) and write $\mathcal{F}$ in these terms. Let us introduce the two kinds of coupling constants T_A : even τ_μ for odd operators $[\Phi_\mu, G_-]$ and odd θ_μ for even Φ_μ .

The main object in the commutativity equations in terms of $\mathcal{F}$ is

$$B_\mu = \frac{\partial \mathcal{F}}{\partial \theta_\mu} \Big|_{\theta=0}\tag{36}$$

In fact, one can show that (34) leads to

$$\begin{aligned}\mathcal{F}|_{\theta=0} &= 0 \quad , \quad \text{hence} \\ \frac{\partial}{\partial \theta_\mu} \frac{\partial}{\partial \theta_\nu} \{\mathcal{F}, \mathcal{F}\}|_{\theta=0} &= [\partial_\mu \mathcal{F}|_{\theta=0}, \partial_\nu \mathcal{F}|_{\theta=0}] = 0\end{aligned}\tag{37}$$

The equation (37) would be the commutativity equation if one demonstrates that $B_\mu = \frac{\partial B}{\partial \tau_\mu}$. It follows from the properties of G_-

$$\begin{aligned}B_\mu &= \frac{\partial B}{\partial \tau_\mu} \implies \frac{\partial B_\nu}{\partial \tau_\mu} = \frac{\partial B_\mu}{\partial \tau_\nu} \sim \\ ..\Phi_\mu G_+ e^{-tH} [\Phi_\nu, G_-] ... &= ..[\Phi_\mu, G_-] G_+ e^{-tH} \Phi_\nu ... \end{aligned}\tag{38}$$

6. Consider the deformation of solutions to the anticommutativity equation by the variation of operator G_+ in terms of $\mathcal{K} = \frac{G_+}{H} = \int_0^\infty G_+ e^{-tH} dt$ so that

$$\{\mathcal{K}, Q\} = 1 - \Pi_0\tag{39}$$

[§]This equation is contained as the t -part in the $t - t^*$ equations of [5].

The variation of $\mathcal{K}$ satisfies

$$\{\delta\mathcal{K}, Q\} = 0 \implies \delta\mathcal{K} = [Q, Z] \quad (40)$$

The variation $\delta\mathcal{K}$ is exact, since Q does not have cohomologies in V_1 . We can write the variation of the solution to the anticommutativity equation and after some algebra we get

$$\delta_{\mathcal{K}}\mathcal{F} = -\{\mathcal{D} - \mathcal{F}, \mathcal{F}_Z\}_s \quad (41)$$

In (41) $\mathcal{F}_Z$ is obtained from the expression[¶] for $\mathcal{F}$ in terms of $\mathcal{K}$ by replacing insertions of $\mathcal{K}$ by Z . The variation of $\mathcal{F}$ in the form (41) is a counterpart of the gauge transformations in the zero curvature equation (24) because they retain this equation

$$\begin{aligned} \{\mathcal{D} - \mathcal{F}, \mathcal{D} - \mathcal{F}\} &= 0 \implies \{\mathcal{D} - \mathcal{F}, \delta\mathcal{F}\} = 0 \\ \{\mathcal{D} - \mathcal{F}, \delta_{\mathcal{K}}\mathcal{F}\} &= \{\mathcal{D} - \mathcal{F}, \{\mathcal{D} - \mathcal{F}, \mathcal{F}_Z\}_s\} = 0 \end{aligned} \quad (42)$$

The special property of commutativity equation is that B_μ is invariant under the variations of $\mathcal{K}$, it follows from (41) and (34). Actually $\mathcal{F}|_{\theta=0} = 0$ and $\mathcal{F}_Z|_{\theta=0} = 0$, hence the variation term linear in θ is

$$\delta_{\mathcal{K}}B_\mu = \delta_{\mathcal{K}}\frac{\partial\mathcal{F}}{\partial\theta_\mu}|_{\theta=0} = 0 \quad (43)$$

In the case of anticommutativity equation one generally has only two terms invariant under the variations of $\mathcal{K}$: $\mathcal{F}^{(1)}$ and $\mathcal{F}^{(2)}$.

Acknowledgements: The author is grateful to A. Morozov, A. Mironov and especially to A. Losev for fruitful discussions. The author also acknowledges the kind hospitality in Kaiserslautern University during the preparation of this work. This work is partly supported by grant RFBR 01-02-17682-a and by the Volkswagen-Stiftung.

References:

- [1] E. Witten, "On the structure of the topological phase of two-dimensional gravity", Nucl.Phys.B340:281-332, 1990
- [2] E. Witten, "Quantum Field Theory and the Jones Polynomial", Commun.Math.Phys.121:351,1989
- [3] E. Witten, "Topological quantum field theories", Commu.Math.Phys.117:353,1988
- [4] E. Witten, "Topological sigma models", Commun.Math.Phys.118:411,1988
- [5] S. Cecotti, C. Vafa, "Topological antitopological fusion", Nucl.Phys.B367:359-461,1991
- [6] A. Losev, "Commutativity equations, operator-valued cohomologies of the "sausage" compactification of $(C^*)^N/C^*$ and SQM" Preprint ITEP-TH-84/98, LPTHE-61/98.
- [7] A. Losev and Yu. Manin, "New moduli space of pointed curves and family of flat connections", AG/0001003
- [8] A. Losev and I. Polyubin, "On compatibility of tensor products on solutions to commutativity and WDVV equations", Pisma Zh.Eksp.Teor.Fiz.73:59-63,2001

[¶] Schematically, $\mathcal{F} = \sum \Pi_0 \Phi \mathcal{K} \Phi \mathcal{K} \dots \Phi \mathcal{K} \Phi \Pi_0$.